



Sea level anomalies control phytoplankton biomass in the Costa Rica Dome area

M. Kahru,¹ P. C. Fiedler,² S. T. Gille,¹ M. Manzano,³ and B. G. Mitchell¹

Received 15 August 2007; revised 18 September 2007; accepted 17 October 2007; published 17 November 2007.

[1] Satellite data show that chlorophyll-a concentration (Chl-a) in the northeastern tropical Pacific is well correlated with sea level anomaly (SLA). This correlation spans a wide spectrum of scales from large-scale phenomena like ENSO to mesoscale cyclonic and anticyclonic eddies. Negative SLA (e.g. during La Niña events and in cyclonic eddies) is associated with the lifting of isopycnals in the nutricline and increased Chl-a due to enhanced phytoplankton growth, while positive SLA (e.g. during El Niño events and in anticyclonic eddies) is associated with a deeper nutricline and reduced Chl-a due to decreased phytoplankton growth. The coupling between SLA and Chl-a anomaly in the Costa Rica Dome (CRD) area is tighter than has previously been recorded anywhere in the world ocean. 70% of the interannual variations in Chl-a anomaly in the CRD area is explained by a combination of the positive and negative effects of SLA. **Citation:** Kahru, M., P. C. Fiedler, S. T. Gille, M. Manzano, and B. G. Mitchell (2007), Sea level anomalies control phytoplankton biomass in the Costa Rica Dome area, *Geophys. Res. Lett.*, 34, L22601, doi:10.1029/2007GL031631.

1. Introduction

[2] Wind jets blowing through mountain gaps of southern Mexico and Central America into the Pacific Ocean are known to produce a rapid oceanic response leading to the generation of oceanic eddies in the Gulfs of Tehuantepec, Papagayo and Panama [Stumpf, 1975; Stumpf and Legeckis, 1977; McCreary *et al.*, 1989; Barton *et al.*, 1993; Chelton *et al.*, 2000; Willett *et al.*, 2006]. These eddies have long been suspected to be important in transporting nutrients and biogenic material from the continental margin to the interior of the ocean, a process recently quantified by Samuelsen [2005]. Indeed, the eddies are known to have distinctive signatures in ocean color [Müller-Karger and Fuentes-Yaco, 2000; Gonzalez-Silvera *et al.*, 2004] and sea-surface height [Willett, 1996]. While most studies have dealt only with the anticyclonic eddies which are larger and have longer lifespan, the cyclonic eddies may be as numerous but dissipate relatively quickly [Gonzalez-Silvera *et al.*, 2004]. Palacios and Bograd [2005] performed a census of eddies and found that the weaker cyclonic eddies did not satisfy their criteria of eddy detection. They also found that the anticyclonic eddy activity (the number of anticyclonic eddies, their

intensity and lifespan) intensified during El Niño years and decreased during La Niña years.

[3] ‘Eddy pumping’ has been suggested as a mechanism by which ocean eddies and meanders on scales of tens to hundreds of kilometers drive the vertical motion of water, including lifting subsurface nutrients into the euphotic zone [Falkowski *et al.*, 1991; McGillicuddy *et al.*, 1998; Oschlies and Garçon, 1998]. Wilson and Adamec [2001, 2002] and Wilson and Coales [2005] have analyzed global relationships between sea surface height, mixed layer depth, thermocline depth and Chl-a and concluded that the dominant mechanism in the tropics is the thermocline uplift corresponding to negative sea level anomaly that enhances phytoplankton growth and concentration. With the availability of high-quality, long time series of sea level anomaly [Ducret *et al.*, 2000] and ocean color [McClain *et al.*, 2006] it is now possible to map important relationships of the phytoplankton environment at scales of tens and hundreds of kilometers using satellite data.

2. Data and Methods

[4] Satellite-derived Level-2 and Level-3 datasets of chlorophyll-a concentration (Chl-a) were obtained from the NASA Ocean Color website (<http://oceancolor.gsfc.nasa.gov/>). Monthly and 8-day Chl-a composites calculated from OCTS (1996–1997) and SeaWiFS (1997–2007) normalized water-leaving radiances using the maximum band-ratio algorithm [O’Reilly *et al.*, 1998] were used to create monthly and 8-day anomalies as the ratio of a current Chl-a to the long-term average Chl-a of the same time period and expressed as percentage anomaly, i.e. $100 \times (\text{Anomaly} - 1)$. Similarly, sea-surface temperature (SST) anomalies were constructed from the monthly Pathfinder version 5 data (<http://www.nodc.noaa.gov>) and expressed as the difference from the long-term mean SST of the same period.

[5] Weekly maps of SLA merged from TOPEX/POSEIDON, Jason and ERS-1/2 data created by AVISO [Ducret *et al.*, 2000] for the 1992–2006 period were used (<http://www.aviso.oceanobs.com>). As the detection of individual eddies by the shape of the SLA field, e.g. by the existence of certain closed contours, is rather subjective, we do not try to separate individual eddies here but attempt to measure eddy activity in general. As a measure of cyclonic eddy activity we calculated monthly root mean square (RMS) of negative SLA in 7×7 pixel windows (RMSNEG) of the updated SLA dataset. Similarly, monthly root mean square of positive SLA in 7×7 pixel windows (RMSPOS) was used as a measure of the anticyclonic eddy activity. Monthly time series of SLA, RMSNEG and RMSPOS were correlated with monthly Chl-a anomalies.

¹Scripps Institution of Oceanography, University of California, San Diego, La Jolla, California, USA.

²Southwest Fisheries Science Center, NOAA National Marine Fisheries Service, La Jolla, California, USA.

³Centro de Investigaciones Biológicas del Noroeste, La Paz, Baja California Sur, Mexico.




Figure 1. Correlation coefficient r of Chl-a anomaly with (a) sea level anomaly (SLA), (b) RMS of negative SLA, and (c) RMS of positive SLA. Isolines for r equal to $|0.25|$ and $|0.5|$ are shown. The spatial domain CRD ($\sim 720 \times 330$ km) in Figure 1b specified as an area with $r > 0.5$ is the approximate location of the Costa Rica Dome. $|r| > 0.23$ is significant at $P < 0.01$.

[6] To evaluate the correlation between two time series at different frequency bands we used squared coherency. Squared coherency was calculated with STATISTICA™ v.6 (<http://www.statsoft.com>) and is considered significant if it exceeds

$$\frac{2F_{2,\nu-2}}{\nu-2+2F_{2,\nu-2}} = 1 - \alpha^{\frac{2}{\nu-2}}$$

This relationship is derived from the F -distribution with $\alpha = 0.05$ and ν being the number of equivalent degrees of freedom. Bias in the squared coherency is related to spectral estimator and the number of ν . Small number of ν may cause a large positive bias in coherency estimates, whereas low frequency peaks may be affected by over-smoothing (see *von Storch and Zwiers* [1999] for further details). For monthly time series we found that squared coherency peaks higher than 0.39 are significant ($P < 0.05$; significance estimated for $\nu = 14$).

3. Results

[7] Figure 1a shows the spatial distribution of the correlation coefficient between Chl-a anomaly and SLA. Chl-a anomaly has significant negative correlation with SLA over large areas of northeastern tropical Pacific, most notably downstream of the typical path of the eddy trajectories [e.g., *Palacios and Bograd*, 2005]. The negative correlation becomes insignificant north of about 10° N latitude, where the thermocline deepens [*Fiedler*, 2002, Figure 1]. Spatial correlation patterns between Chl-a anomaly and the RMS of negative SLA and the RMS of positive SLA (Figures 1b and 1c) are approximately similar but with opposite sign. RMSNEG can be considered a proxy for cyclonic eddy activity, and RMSPOS is a proxy for anticyclonic eddy activity. The compact area of strong positive correlation between RMSNEG and Chl-a anomaly (~ 720 km east-west \times 330 km north-south) defined by $r \geq 0.5$ (CRD in Figure 1b) coincides with the general location of the Costa Rica Dome, an area with a shallow, doming thermocline [*Fiedler*, 2002]. Time series of various parameters were constructed in this spatial domain (Figures 2 and 3).

[8] The response in Chl-a to the drop in negative SLA (Figure 3, bottom) and a sharp increase in positive SLA (Figure 4, bottom) during the 1997–1998 El Niño event was practically instantaneous and no delay in the response of Chl-a could be detected. Even when comparing 8-day Chl-a and 7-day SLA data (not shown) no significant delay could be detected. The El Niño event of 1997–1998 was extremely anomalous and may cause the high correlations between SLA and Chl-a anomaly. We therefore excluded the drastic El Niño event in the calculations and repeated the calculations for the 1999–2006 period but the correlations




Figure 2. (top) Scatter plot and (bottom) time series of monthly mean SLA (black line, left axis) and Chl-a anomaly (gray line, right axis) in area CRD of Figure 1b.




Figure 3. (top) Scatter plot and (bottom) time series of monthly mean RMS of negative SLA (black line, left axis) and Chl-a anomaly (gray line, right axis) in area CRD of Figure 1b.

stayed highly significant, although their absolute values decreased (Table 1). For example, r^2 between RMS of negative SLA decreased from 0.663 for the whole period to 0.563 for the 1999–2006 period. We can predict Chl-a anomaly in the Costa Rica Dome area with the following equation:

$$\text{ChlAnom} = 0.789 + 0.051 * \text{RMSNEG} - 0.017 * \text{RMSPOS}, \quad (1)$$

where RMSNEG and RMSPOS are the RMS of negative and positive SLA, respectively. This relationship explains over 70% of the total variance ($R^2 = 0.705$) in Chl-a anomaly. The positive relationship between SLA and SST anomaly (not shown) is even stronger ($r^2 = 0.725$ for the 1999–2006 period).

[9] Visual inspection of Figures 2, 3, and 4 shows that in addition to the strong correlation between SLA and Chl-a anomaly at large temporal scales, e.g. El Niño-Southern Oscillation (ENSO), many features at much shorter scales are also correlated. We confirmed this with cross-spectral analysis. Figure 5 shows that the squared coherence (similar to the squared coefficient of correlation for certain frequencies) shows significant correlation at a number of time scales. The significant coherence at 2.4 months is likely to be caused by the passage of mesoscale eddies. The high coherence at longest scales is consistent with ENSO vari-




Figure 4. (top) Scatter plot and (bottom) time series of monthly mean RMS of positive SLA (black line, left axis) and Chl-a anomaly (gray line, right axis) in area CRD of Figure 1b.

ability. The intermediate scales of high coherence may be due to seasonal changes.

4. Discussion

[10] The negative correlation between Chl-a and SLA, especially in the tropics, has been observed earlier [Wilson and Adamec, 2001] and explained by deepening of the thermocline (and associated nutricline) at positive SLA and shoaling of the thermocline (and nutricline) at negative SLA. The increase in phytoplankton biomass (Chl-a) in cyclonic eddies is believed to be the result of increased local production due to the lifting of the high-nutrient waters into the euphotic layer in areas of shoaling of the thermocline [Benitez-Nelson *et al.*, 2007]. The interactions between Chl-

Table 1. Coefficient of Determination Between SLA and Its Positive and Negative Parts With Chl-a Anomaly in Area CRD of Figure 1b^a

Variable	1996–2006	1999–2006
SLA	0.663	0.553
RMS of negative SLA	0.643	0.572
RMS of positive SLA	0.490	0.314
RMSNEG & RMSPOS	0.705	0.579

^aCoefficient of determination, r^2 . “RMSNEG & RMSPOS” means using both RMS of negative and positive SLA in a multiple linear regression to predict Chl-a anomaly. Significance was tested for the “effective” degrees of freedom caused by autocorrelation and results were significant in all cases ($P < 0.01$).




Figure 5. Squared coherency spectrum between monthly time series of RMS of negative SLA (RMSNEG) and Chl-a anomaly in the Costa Rica Dome area. The peaks at frequencies 0.19, 0.28, 0.41 month⁻¹ correspond to periods of 5.3, 3.6, and 2.4 months, respectively. The dashed horizontal line indicates the approximate 95% significance level.

a and eddy dynamics are very complex and involve a number of mechanisms which are not well known. Upwelling at the periphery of anticyclonic eddies [McGillicuddy *et al.*, 2007; Kahru *et al.*, 2007] can cause local increase of Chl-a. However, in our satellite study with ~25 km resolution of SLA and 9-km resolution of Chl-a we cannot effectively evaluate these small-scale mechanisms and have to resort to statistical correlations at relatively low spatial scales. Contrary to Lopez-Calderon *et al.* [2006] who concluded that both cyclonic and anticyclonic eddies increase local Chl-a in the north-eastern tropical Pacific, we find significant negative correlation between Chl-a and positive SLA, a proxy for anticyclonic eddies. Both cyclonic and anticyclonic eddies often occur in groups and when using low-resolution, average SLA, the negative and positive anomalies can partly cancel each other and their net effect is altered. When the positive and negative components of SLA are separated then the negative effects of the positive SLA are obvious (Figure 4). It is also significant that when using the positive and negative components of SLA separately, the resulting coefficient of determination is higher than when using the mean SLA alone (0.705 versus 0.663 for the whole period). This may be caused by the smoothing effect of the 7×7 pixel window when calculating the RMS of SLA. Cross-spectral analysis confirms that Chl-a anomaly is coherent with SLA at a number of temporal scales, e.g. corresponding to ENSO, seasonal changes and mesoscale eddies.

[11] An alternative hypothesis (the “hay rake” mechanism) for the apparent increase in Chl-a was proposed by Dandonneau *et al.* [2003] who argued that ocean color anomalies associated with sea level anomalies are not actually caused by increased Chl-a but rather by the accumulation of detrital material in convergence zones of ocean eddy fields. The “hay rake” mechanism does not seem to be valid in the Costa Rica Dome area as periods of increased Chl-a are associated with periods of cold SST that is consistent with upwelling of cold, nutrient-rich waters due to “eddy pumping”.

[12] McGillicuddy *et al.* [2007] recently proposed a new mechanism for sustained upwelling in anticyclonic mode-

water eddies caused by eddy-wind interactions. The eddy/wind interaction tends to sustain anticyclonic eddies and dampen cyclonic eddies. This may be one of the explanations of the longer life time of anticyclonic eddies. However, in this region increased Chl-a is clearly associated with cyclonic eddies and eddy/wind interaction is expected to reduce the effect of eddy-induced upwelling.

5. Conclusions

[13] The coupling between sea level anomaly and Chl-a anomaly in the Costa Rica Dome area is tighter than has previously been reported anywhere in the global ocean. Negative SLA is associated with increased Chl-a, and positive SLA with decreased Chl-a. Positive or negative SLA reflects, respectively, deepening or shoaling of the thermocline. This mechanism is especially effective in modulating the nutrient supply to the euphotic zone in the Costa Rica Dome area due to its shallow thermocline, close proximity of the thermocline and the nutricline, and the effect of the energetic mesoscale eddies impinging on the area. 70% of the interannual variations in Chl-a anomaly in the Costa Rica Dome area is explained by the combination of the positive effects of the negative SLA (including La Niña and cyclonic eddies) and the negative effects of the positive SLA (including El Niño and anticyclonic eddies).

[14] **Acknowledgments.** Financial support to MK and BGM was provided by the NASA Ocean Biology and Biogeochemistry Program (NNG05GR23G) and the NSF (OPP grant ANT04-44134 and OCE grant 04-17616). STG received support from NASA JPL contract 1224031. We thank the NASA Ocean Color Processing Group and AVISO for satellite data. This is a CCE LTER contribution 0050.

References

- Barton, E. D., M. L. Argote, J. Brown, P. M. Kosro, M. Lavin, J. M. Robles, R. L. Smith, A. Trasvina, and H. S. Velez (1993), Supersquirt: Dynamics of the Gulf of Tehuantepec, Mexico, *Oceanography*, 6, 23–30.
- Benitez-Nelson, C. R., et al. (2007), Mesoscale eddies drive increased silica export in the subtropical Pacific Ocean, *Science*, 316, 1017–1021, doi:10.1126/science.1136221.
- Chelton, D. B., M. H. Freilich, and S. K. Esbensen (2000), Satellite observations of the wind jets off the Pacific coast of Central America. Part I: Case studies and statistical characteristics, *Mon. Weather Rev.*, 128, 1993–2018.
- Dandonneau, Y., A. Vega, H. Loisel, Y. du Penhoat, and C. Menkes (2003), Oceanic Rossby waves acting as a “hay rake” for ecosystem floating by-products, *Science*, 302, 1548–1551.
- Ducet, N., P.-Y. Le Traon, and G. Reverdin (2000), Global high resolution mapping of ocean circulation from TOPEX/Poseidon and ERS-1/2, *J. Geophys. Res.*, 105, 19,477–19,498.
- Falkowski, P. G., D. Ziemann, Z. Kolber, and P. K. Bienfang (1991), Role of eddy pumping in enhancing primary production, *Nature*, 352, 55–58.
- Fiedler, P. C. (2002), The annual cycle and biological effects of the Costa Rica Dome, *Deep Sea Res., Part I*, 49, 321–338.
- Gonzalez-Silvera, A., E. Santamaria-del-Angel, R. Millán-Nuñez, and H. Manzo-Monroy (2004), Satellite observations of mesoscale eddies in the Gulfs of Tehuantepec and Papagayo (eastern tropical Pacific), *Deep Sea Res., Part II*, 51, 587–600.
- Kahru, M., B. G. Mitchell, S. T. Gille, C. D. Hewes, and O. Holm-Hansen (2007), Eddies enhance biological production in the Weddell-Scotia Confluence of the Southern Ocean, *Geophys. Res. Lett.*, 34, L14603, doi:10.1029/2007GL030430.
- Lopez-Calderon, J., H. Manzo-Monroy, E. Santamaria-del-Angel, R. Castro, A. Gonzalez-Silvera, and R. Millán-Nuñez (2006), Mesoscale variability of the Mexican Tropical Pacific using TOPEX and SeaWiFS data, *Cienc. Mar.*, 32, 539–549.
- McClain, C., S. Hooker, G. Feldman, and P. Bontempi (2006), Satellite data for ocean biology, biogeochemistry, and climate research, *Eos Trans. AGU*, 87, 337.
- McCreary, J. P., Jr., H. S. Lee, and D. B. Enfield (1989), The response of the coastal ocean to strong offshore winds: With application to

- circulations in the gulfs of Tehuantepec and Papagayo, *J. Mar. Res.*, *47*, 81–109.
- McGillicuddy, D. J., A. R. Robinson, D. A. Siegel, H. W. Jannasch, R. Johnson, T. D. Dickey, J. McNeil, A. F. Michaels, and A. H. Knap (1998), Influence of mesoscale eddies on new production in the Sargasso Sea, *Nature*, *394*, 263–266.
- McGillicuddy, D. J., et al. (2007), Eddy/wind interactions stimulate extraordinary mid-ocean plankton blooms, *Science*, *316*, 1021–1026, doi:10.1126/science.1136256.
- Müller-Karger, F. E., and C. Fuentes-Yaco (2000), Characteristics of wind-generated rings in the eastern tropical Pacific Ocean, *J. Geophys. Res.*, *105*, 1271–1284.
- O'Reilly, J. E., S. Maritorena, B. G. Mitchell, D. A. Siegel, K. L. Carder, S. A. Garver, M. Kahru, and C. R. McClain (1998), Ocean color chlorophyll algorithms for SeaWiFS, *J. Geophys. Res.*, *103*, 24,937–24,953.
- Oschlies, A., and V. Garçon (1998), Eddy-induced enhancement of primary production in a model of the north Atlantic Ocean, *Nature*, *394*, 266–269.
- Palacios, D. M., and S. J. Bograd (2005), A census of Tehuantepec and Papagayo eddies in the northeastern tropical Pacific, *Geophys. Res. Lett.*, *32*, L23606, doi:10.1029/2005GL024324.
- Samuelson, A. (2005), Modeling the effect of eddies and advection on the lower trophic ecosystem in the northeast tropical Pacific, Ph.D. thesis, 86 pp., Florida State Univ., Tallahassee.
- Stumpf, H. G. (1975), Satellite detection of upwelling in the Gulf of Tehuantepec, Mexico, *J. Phys. Oceanogr.*, *5*, 648–658.
- Stumpf, H. G., and R. V. Legeckis (1977), Satellite observations of mesoscale eddy dynamics in the eastern tropical Pacific Ocean, *J. Phys. Oceanogr.*, *7*, 648–658.
- von Storch, H., and F. W. Zwiers (1999), *Statistical Analysis in Climate Research*, 484 pp., Cambridge Univ. Press, New York.
- Wilson, C., and D. Adamec (2001), Correlations between surface chlorophyll and sea surface height in the tropical Pacific during the 1997/1999 El Niño-Southern Oscillation event, *J. Geophys. Res.*, *106*, 31,175–31,188.
- Wilson, C., and D. Adamec (2002), A global view of bio-physical coupling from SeaWiFS and TOPEX satellite data, 1997–2001, *Geophys. Res. Lett.*, *29*(8), 1257, doi:10.1029/2001GL014063.
- Wilson, C., and V. J. Coales (2005), Global climatological relationships between satellite biological and physical observations and upper ocean properties, *J. Geophys. Res.*, *110*, C10001, doi:10.1029/2004JC002724.
- Willett, C. S. (1996), A study of anticyclonic eddies in the eastern tropical Pacific Ocean with integrated satellite, in situ, and modeled data, Ph.D. thesis, 127 pp., Univ. of Colo., Boulder.
- Willett, C. S., R. Leben, and M. F. Lavin (2006), Eddies and mesoscale processes in the eastern tropical Pacific: A review, *Prog. Oceanogr.*, *69*, 218–238.

P. C. Fiedler, Southwest Fisheries Science Center, NOAA National Marine Fisheries Service, La Jolla, CA 92037, USA.

S. T. Gille, M. Kahru, and B. G. Mitchell, Scripps Institution of Oceanography, University of California, San Diego, 9500 Gilman Drive, La Jolla, CA 92093-0218, USA. (mkahru@ucsd.edu)

M. Manzano, Centro de Investigaciones Biológicas del Noroeste, La Paz, BCS 23090, México.